

Pubic Symphysis

Suchey Brooks (1988), and Todd (1920)

Traits	Left	Right
Syphyseal Face	V	
Syphyseal Rim	V	
Upper Extremity	X	
Lower Extremity	X	
Dorsal Margin	IV	
Ventral Margin	X	
Other Features	X	
Phase/Age	Phase: V Age: 25- 74 Mean: 49	

Auricular surface

Lovejoy et al. (1985), and Buckberry and Chamberlain (2002)

Traits	Left	Right
Transverse organisation	5	5
Surface texture	4	4
Micro-porosity	3	3
Macro-porosity	2	2
Apical changes	3	3
Phase/Age	Phase: VII Age: 53-92 Mean: 72	Phase: VII Age: 53-92 Mean: 72

DATA FYSISCHE ANTROPOLOGIE

Complex method for four or less age indicators

Broca (1875), Nemeskéri, et. al. (1960), Acsádi and Nemeskéri (1970), Sjøvold (1975), and WEA (1980)

Age indicator	Phase
Sympyseal face of pubis of males	V
Females only if without parturition scars	
Spongiosa of proximal femur	
Spongiosa of proximal humerus	
Endocranial suture obliteration	

Skeletal age	Age (years)	Range (years)	Interval (years)	Confidence
In case of 4 ages indicators		± 2.5 years	-	80%-85%
In case of 3 ages indicators		± 3.0 years	-	80%-85%
In case of 2 ages indicators			-	80%-85%
In case of 1 age indicator			60-80	/

Final age assessment	53-74 oud volwassen
----------------------	------------------------

Notes: