

DATA FYSISCH ANTROPOLOGIE

Pubic Symphysis Suchey Brooks (1988), and Todd (1920)		
Traits	Left	Right
Symphyseal Face	4	4
Symphyseal Rim	4	4
Upper Extremity	4	4
Lower Extremity	3	3
Dorsal Margin	5	5
Ventral Margin	3	3
Other Features		
Phase/Age	Phase IV Mean 35,2 Range 23-57	Phase IV Mean 35,2 Range 23-57

Auricular surface Lovejoy et al. (1985), and Buckberry and Chamberlain (2002)		
Traits	Left	Right
Transverse organisation	3	
Surface texture	2	
Micro-porosity	3	
Macro-porosity	2	
Apical changes	1	
Phase/Age	Phase IV Mean 51 Range 29-81	

DATA FYSISCH ANTROPOLOGIE

Complex method for four or less age indicators

Broca (1875), Nemeskéri, et. al. (1960), Acsádi and Nemeskéri (1970), Sjøvold (1975), and WEA (1980)

Age indicator			Phase	
Symphyseal face of pubis of males Females only if without parturition scars			IV	
Spongiosa of proximal femur				
Spongiosa of proximal humerus				
Endocranial suture obliteration				
Skeletal age	Age (years)	Range (years)	Interval (years)	Confidence
In case of 4 ages indicators		± 2.5 years	-	80%-85%
In case of 3 ages indicators		± 3.0 years	-	80%-85%
In case of 2 ages indicators			-	80%-85%
In case of 1 age indicator			50-70	/

Ossification at the sternal end of the 4th rib

Işcan and Loth (1984, 1985)

Phase III	Age interval in years 95% confidence
24 - 28	-

Final age assessment

29-57
Midden volwassen