

DATA FYSISCH ANTROPOLOGIE

Pubic Symphysis Suchey Brooks (1988), and Todd (1920)		
Traits	Left	Right
Symphyseal Face	4	5
Symphyseal Rim	4	5
Upper Extremity	4	3
Lower Extremity	4	4
Dorsal Margin	5	4
Ventral Margin	5	4
Other Features	5	5
Phase/Age	Phase v Mean 45,6 Range 27-66	Phase v Mean 45,6 Range 27-66

Auricular surface Lovejoy et al. (1985), and Buckberry and Chamberlain (2002)		
Traits	Left	Right
Transverse organisation	5	5
Surface texture	3	3
Micro-porosity	3	3
Macro-porosity	2	2
Apical changes	3	3
Phase/Age	Phase VI Mean 67 Range 39-91	Phase VI Mean 67 Range 39-91

DATA FYSISCH ANTROPOLOGIE

Ectocranial suture closure Meindle and Lovejoy (1985)			
	Suture sector	Score (0 – 3)	
1	Midlambdoid	1	
2	Lambda	2	
3	Obelion	3	
4	Antior sagittal	2	
5	Bregma	2	
6	Midcoronal	1	
7	Pterion	2	
8	Sphenofrontal	3	
9	Inferior sphenotemporal	1	
10	Superior sphenotemporal	3	
Vault composite (1-7) 13		Mean	SD
		45,2	12,6
Lateral-anterior composite (6-10) 10		Mean	SD
		51,9	12,5
		Range	
		24-75	
		Range	
		33-76	

Complex method for four or less age indicators Broca (1875), Nemeskéri, et. al. (1960), Acsádi and Nemeskéri (1970), Sjøvold (1975), and WEA (1980)				
Age indicator			Phase	
Symphyseal face of pubis of males Females only if without parturition scars			V	
Spongiosa of proximal femur				
Spongiosa of proximal humerus				
Endocranial suture obliteration				
Skeletal age	Age (years)	Range (years)	Interval (years)	Confidence
In case of 4 ages indicators		± 2.5 years	-	80%-85%
In case of 3 ages indicators		± 3.0 years	-	80%-85%
In case of 2 ages indicators			-	80%-85%
In case of 1 age indicator			60-80	/

DATA FYSISCHE ANTROPOLOGIE

Final age assessment	39-76 Midden volwassen
-----------------------------	---------------------------

Notes: